FUNCTIONAL ANALYSIS & FUNCTIONAL LIE ALGEBRAS

RIGOROUS MATHS IN QUANTUM

Rigorous Operator Foundations, Infinite Dimensions, and Haar Measures

An axiomatic mathematical guide detailing the rigorous functional analytic structures that prevent quantum paradoxes.

1. THE BOUNDEDNESS PARADOX & UNBOUNDED OPERATORS

In finite-dimensional quantum mechanics, every linear operator is bounded. However, when working in infinite-dimensional Hilbert spaces ($L^2(\mathbb{R})$), physical observables like position (X) and momentum (P) are natively **unbounded operators**. This creates severe domains of definition issues.

[infobox] **The Hellinger-Toeplitz Constraint:** A standard mathematical theorem states that any symmetric operator defined on the entire Hilbert space must be bounded. Because position and momentum are unbounded, they cannot be defined everywhere in $\mathcal{H}$:

$$X\psi(x) = x\psi(x) \qquad \text{and} \qquad P\psi(x) = -i\hbar\frac{d\psi(x)}{dx}$$

The Consequence: The Heisenberg commutation relation $[X, P] = i\hbar I$ can **never** exist as a finite-dimensional matrix relation. Taking the matrix trace reveals $\mathrm{Tr}(XP - PX) = 0$, which contradicts $\mathrm{Tr}(i\hbar I) = i\hbar \cdot d$. Infinite dimensions are an analytical absolute.

2. THE MATRIX ILLUSION: SYMMETRIC VS. SELF-ADJOINT OPERATORS

In standard text treatments, the terms "Hermitian" and "Self-Adjoint" are thrown around interchangeably. For infinite dimensions, this mathematical looseness leads to catastrophic physical paradoxes (such as complex eigenvalues for energy).

The Domain Distinction: Let A be an operator with a domain $\mathcal{D}(A)$, and its adjoint $A^\dagger$ have a domain $\mathcal{D}(A^\dagger)$.

b✦

leftmargin=*, noitemsep, topsep=2ptSymmetric (Hermitian): $\langle\phi|A\psi\rangle = \langle A\phi|\psi\rangle$ for all $\phi, \psi \in \mathcal{D}(A)$. This only demands $\mathcal{D}(A) \subseteq \mathcal{D}(A^\dagger)$.

Self-Adjoint: Satisfies the symmetric condition **and** requires their structural operational domains to be perfectly identical: $\mathcal{D}(\mathbf{A}) = \mathcal{D}(\mathbf{A}^\dagger)$.

The Mathematical Pitfall: Only true self-adjoint operators satisfy the Spectral Theorem, guaranteeing exclusively real eigenvalues and deterministic unitary time evolution via Stone's Theorem ($U = e^{-iHt/\hbar}$).

3. CONTINUUMS GELFAND TRIADS (RIGGED HILBERT SPACES)

Dirac notation frequently references continuous states like plane waves $|p\rangle$ or delta functions $|x\rangle$. Rigorously, these vectors do not exist inside a standard Hilbert space $\mathcal{H}$ because their norm is infinite ($\langle x|x\rangle = \delta(0) = \infty$), violating Postulate 1.

THE RIGGED HILBERT SPACE (GELFAND TRIAD)

To reconcile Dirac's calculation structures with pure functional analysis, the vector layout is wrapped into a nested mathematical sandwich:

$$\mathbf{\Phi} \subset \mathcal{H} \subset \mathbf{\Phi}^*$$

The Components: Φ is the space of well-behaved test functions (e.g., Schwartz space), $\mathcal{H}$ is the physical L^2 space, and Φ^* is the topological dual containing generalized distributions. Continuous states exist cleanly within Φ^*, preserving operational validity.

4. INTEGRATION ON TOPOLOGICAL GROUPS: THE HAAR MEASURE

When proving theorems about random quantum circuits, compiling gates, or measuring variational properties in QML, algorithms must evaluate averages over entire continuous gate groups (such as $SU(2^N)$). This requires an entry-wise invariant metric—the **Haar Measure** $d\mu$.

The Invariance Invariant: For any arbitrary elements V belonging to the compact Lie group G, the Haar probability distribution guarantees that shifting the entire operator geometry preserves total area metrics:

$$\int_G f(U)d\mu(U) = \int_G f(VU)d\mu(U) = \int_G f(UV)d\mu(U) \qquad \text{where } \int_G d\mu(U) = 1$$

The Application Link: The statistical framework of Haar integration enables the rigorous proofs behind the **Barren Plateau paradox** and evaluates the complexity limits of random quantum state t-designs.